


Arsenic in Soils, Mine Tailings, and Former Industrial Sites

Guillaume Morin and Georges Calas¹



SEM micrograph of sediments from the Carnoules (France) acid mine drainage site, showing amorphous As^{5+} - Fe^{3+} hydrous oxides associated with bacteria.

MODIFIED FROM MORIN ET AL. (2003)

Much progress has recently been made on the relation between the crystal chemistry of arsenic and its speciation and distribution at the Earth's surface. The investigation of As-impacted soils and acid mine drainages, using synchrotron-based techniques, shows the importance of As adsorption on, or coprecipitation with, hydrous ferric oxides in delaying the long-term impact of As on the biosphere. Arsenic mobility often depends on bacterial activity, with accompanying major seasonal modifications of As speciation, even at extreme As concentrations. Remediation technologies use geochemical affinities between arsenic and specific low-temperature phases to reduce the bioavailability of arsenic.

KEYWORDS: arsenic, contamination, soils, acid mine drainage, speciation

INTRODUCTION

Although arsenic occurs naturally in soils, human activities have greatly increased arsenic contamination in the environment, e.g. through releases associated with mining, waste disposal, indiscriminate use of certain pesticides and herbicides, as well as the manufacture of As-bearing chemicals. Contaminated soils act as a source of arsenic, which can be further dispersed in the environment and may adversely affect the growth of plants and animals and, eventually, human health (Duker et al. 2005). A major source of contamination comes from the breakdown of insoluble As minerals, the arsenides and sulfides. Under the conditions prevailing at the Earth's surface, inorganic arsenic is progressively released as pentavalent arsenate (As^{5+}) or trivalent arsenite (As^{3+}) species, and these form a large variety of minerals (Anthony et al. 2000; O'Day this issue). However, most arsenate and arsenite minerals are not stable in soils over the long term. Arsenic may be encountered in a variety of other forms, such as poorly crystalline As-bearing solids, As-sorbed species, and organic forms of arsenic, mainly methylated compounds and carbohydrate derivatives. The difficulty in providing a robust assessment of As mobility at the Earth's surface comes from the complex interplay among the various forms of As and natural waters, together with the major role exerted by biotic processes (Lloyd and Oremland this issue). Depending on local conditions, As concentration and partitioning between solid phases and solutions may be further modified by changes in pH and redox conditions. For example, the dissolution of As may be counteracted by the precipitation of Fe-oxyhydroxides, which efficiently scavenge As^{3+}

and As^{5+} at neutral pH. The commonly observed increase in solubility of As under reducing conditions is now attributed primarily to the reductive dissolution of Fe^{3+} -oxide phases and the subsequent release of adsorbed As. Aided by a rapidly expanding body of experimental data and by the information provided by modern analytical tools such as synchrotron radiation, a better picture of the control exerted by the crystal chemistry of arsenic on its biogeochemical properties is emerging. With this in mind, we will present some examples of studies of con-

taminated soils, former industrial sites, and mine tailings. These examples show the importance of oxidation and dissolution reactions, microbial activity, and water regime. The investigation of both natural high-arsenic geochemical anomalies and old mining and industrial sites contaminated by arsenic provides good data to establish what the most stable As-bearing phases are.

ARSENIC MINERALOGY AT FORMER INDUSTRIAL SITES

Industrial sites may be contaminated with by-products from the processing of As-bearing ores or by chemicals associated with the arsenic industry. These sites provide examples of major contamination, where arsenic has been mobilized from wastes by meteoric waters and has then reacted with mineral phases present in the environment. At most such sites, interaction of arsenic-rich waters with surrounding soils or bedrock may limit the contamination of the neighboring groundwaters and surface waters.



An example of the latter is provided by an old industrial site located on limestone in southern France (Juillot et al. 1999). For more than a century, sulfuric acid and various arsenic compounds were produced at this site, now under restoration. An integrated study of this contaminated area allowed identification of the source of the arsenic and of secondary As-bearing minerals formed at the site. The latter consist of secondary Ca- and Ca-Mg-arsenates, occurring either as surficial crusts or as transported particles. The arsenates formed as a result of the interaction of the runoff waters with As-bearing wastes and underlying limestone. Key geochemical parameters, such as pH and Ca, As, and sulfate concentrations, were monitored during several field campaigns, together with the proportions of various solid phases present, as determined from Rietveld refinement of X-ray diffraction patterns. It was found that the initially carbonate-buffered waters are strongly acidified (pH = 2.2) and enriched in sulfates as a result of the oxidation of

¹ Institut de Minéralogie et Physique des Milieux Condensés, Université de Paris 6, Université de Paris 7, Institut de Physique du Globe de Paris et CNRS, Case 115, 4 Place Jussieu, 75252 Paris, France
E-mail: morin@impmc.jussieu.fr

wastes containing sulfide ores. The subsequent interaction with the underlying limestone causes the pH to shift back to neutrality, and the concentrations of Ca and Mg also progressively increase. The anhydrous arsenate weillite [Ca(AsO₃OH)] is the first phase formed during the neutralization of acidic As-bearing solutions, followed by pharmacolite (CaHAsO₄•2H₂O). The high stability of these Ca-arsenates at this limestone site may be explained by a common ion effect, in which high Ca activity in the runoff waters inhibits the dissolution of Ca-arsenates. This explains an apparent As concentration three to four times lower in these waters than expected from the solubility product of weillite (Swash and Monhemius 1994). Arsenates such as scorodite (FeAsO₄•2H₂O) are seldom encountered because of the scarcity of iron in the As-rich zones of the site. Field observations on the stability of Ca-arsenates may shed light on their long-term efficiency for remediation in soils, in which As may be rapidly immobilized by treatment with lime slurries, leading to the formation of Ca-hydroxyarsenates (Moon et al. 2004).

MINE SITES AND RELATED ENVIRONMENTS

Mining can release high concentrations of arsenic by means of the oxidation of sulfide minerals, either through the roasting of arsenical ores or through acid mine drainage (AMD) (Williams 2001). The latter results from the atmospheric exposure and weathering of sulfide-bearing rocks, such as the waste rock and mine tailings generated during mining and milling operations. Oxidation of sulfide minerals, often mediated by microbial activity, causes waters to become acidified and enriched in sulfate anions and heavy metals. During this process, an overall two-stage reaction governs the dissolution and further oxidation of As:



The oxidation of sulfide (reaction 1) can be catalyzed directly by bacterial activity and indirectly by the Fe³⁺ resulting from the oxidation of Fe²⁺ by bacteria such as *Acidithiobacillus ferrooxidans*. Bacterial oxidation of dissolved Fe²⁺ also leads to the formation of various iron oxides (Cornell and Schwertmann 2003) and hydroxysulfates (Bigam et al. 1996; Alpers et al. 2000). Arsenite oxidation (reaction 2) is slow, especially under acidic conditions, but may be catalyzed by the activity of bacteria such as *Thiomonas* sp. (Bruneel et al. 2003). Reaction (2) is important because As⁵⁺ is less toxic, less soluble, and adsorbs more efficiently than As³⁺ under acidic conditions.

Trapping of As by iron minerals is an efficient natural attenuation process, which reduces considerably the arsenic concentration in AMD waters. At pH lower than 2.5, As⁵⁺ may be incorporated in jarosite [KFe₃(SO₄)₂(OH)₆] (Savage et al. 2000; Foster et al. 1998). At higher pH, As⁵⁺ can sorb onto iron minerals with high specific surface areas (Fukushi et al. 2003; Courtin-Nomade et al. 2005), such as schwertmannite [Fe₈O₈(OH)₆SO₄] and ferrihydrite (Fe₅HO₈•4H₂O) (for 2.5 ≤ pH ≤ 6 and 6 ≤ pH ≤ 8, respectively; Bigam et al. 1996). Arsenic speciation in mining environments provides an interesting picture of the complex interplay among the sources of the arsenic, the geochemical conditions, the dynamics of the system, and bacterial activity.

One of the first studies of arsenic speciation in mine tailings was that of the Mother Lode district, California, USA (Foster et al. 1998; Brown et al. 1999). A quantitative assessment of As speciation, based on XRD and extended X-ray absorption fine-structure (EXAFS) spectroscopy, was undertaken for three mine wastes: fully oxidized tailings (Ruth mine),

partially oxidized tailings (Argonaut mine), and a roasted sulfide ore (Spenceville mine). As⁵⁺ was the predominant oxidation state in the Ruth and Spenceville mine samples, but mixed oxidation states were observed in the Argonaut mine wastes. These analyses suggested the presence of As⁵⁺ species similar to those found in scorodite and As⁵⁺ adsorbed on goethite (α-FeOOH) and gibbsite [α-Al(OH)₃]. Linear least-squares fits of mine waste EXAFS spectra indicated variable As speciation in each of the three mine wastes. Ferric oxyhydroxides and aluminosilicates (probably clay minerals) bind roughly equal proportions of As⁵⁺ in the Ruth mine. Tailings from the Argonaut mine contain ~20% reduced As bound in arsenopyrite (FeAsS) and arsenical pyrite (FeS_{2-x}As_x) and ~80% As⁵⁺ in a ferric arsenate precipitate similar to scorodite. Roasted sulfide ore from the Spenceville mine contains As⁵⁺, both substituting for sulfate in the crystal structure of jarosite and sorbed to iron-oxide surfaces. These investigations show the importance of the mineralogical composition of the As source material and of sorbed As species.

Recent studies of arsenic-rich mine tailings (Paktunc et al. 2004) suggest that the stability of these mineral phases generally increases with increasing Fe/As ratio. For instance, leaching tests on tailings from the Ketzia River mine, Canada, showed that As adsorbed in small quantities on Fe oxides appears to be less soluble than crystalline minerals such as arseniosiderite [Ca₂Fe₃(AsO₄)₃O₂•3H₂O] (Paktunc et al. 2004). These findings agree with the few available solubility data on crystalline and amorphous As-Fe minerals.

An example of natural attenuation via the formation of As-Fe mineral phases is provided by an AMD at Carnoulès, Gard, France, with exceptionally high concentrations of dissolved arsenite. This AMD is generated by 1.5 million tons of mine tailings containing arsenical pyrite (Leblanc et al. 1996). Here, bacteria play an important role, and this biogeochemical coupling causes spatial and seasonal modifications in As speciation (Morin et al. 2003; Casiot et al. 2005). At or near the source of AMD, the anoxic and acid (pH = 2–3) waters are strongly enriched in Fe (0.5–1 g L⁻¹), As (50–350 mg L⁻¹), and sulfate (1–3 g L⁻¹). Under such conditions, arsenite is the predominant dissolved arsenic species. After 30 m of downflow, 20–60% of the As has been removed via the biogenic precipitation of rare mineral species, such as amorphous ferric arsenic hydroxysulfates and nanocrystalline tooeleite, a ferric arsenite (Fig. 1). This intense microbial activity results in an efficient detoxification of the AMD waters although the As-rich mineral phases formed in the upstream zone are fairly soluble. Further downstream, the progressive decrease of the dissolved arsenic concentration allows the precipitation of schwertmannite, which traps most of the remaining dissolved arsenic. The dissolved arsenic concentration thus decreases to about 1 mg L⁻¹ before the confluence with the non-polluted river, 1.5 km downstream from the waste pile (Casiot et al. 2005). An important finding of the studies carried out on the Carnoulès AMD concerns the influence of bacterial activity on the nature of the minerals forming. In the field, seasonal variation of arsenic speciation has been observed over four years of study in the acidic upstream zone. In winter, only microbial oxidation of Fe²⁺ is efficient, leading to the formation of amorphous Fe³⁺-As³⁺ precipitates together with nanocrystalline tooeleite [Fe₆(AsO₃)₄SO₄(OH)₄•4H₂O] (Fig. 1).

In spring and summer, biotic oxidation of both As³⁺ and Fe²⁺ leads to the precipitation of mixed Fe³⁺-As⁵⁺ hydrous oxides, similar to biominerals occurring at geothermal springs (Inskeep et al. 2004). These distinct mineral phases have been replicated in the laboratory using single bacterial strains isolated from the site. Oxidation of Fe²⁺ by




FIGURE 1 Massive bacterial oxidation of iron and arsenic leads to the formation of spectacular outcrops of arsenic-bearing (up to 20 wt% As) sediments and meter-wide stromatolites (bottom center) at the Carnoules (France) acid mine drainage (**D**). The proportion of seasonally alternating As^{3+} and As^{5+} species associated with ferric oxides can be quantified by XANES spectroscopy (**E**). SEM micrograph of winter sediments (**A**) shows the close association between bacteria and a Fe^{3+} -arsenite mineral, tooeelite. This rare mineral, with a distinctive layer structure (**C**), is the main component of the stromatolite (**B**). Laboratory experiments showed that *Acidithiobacillus ferrooxydans* isolated from the site can form tooeelite by oxidizing Fe^{2+} but not As^{3+} . Modified from Morin et al. (2003)

Acidithiobacillus ferrooxydans strains, which are unable to oxidize As^{3+} , leads to the formation of mixed Fe^{3+} - As^{3+} amorphous hydroxysulfates and nanocrystalline tooeelite (Morin et al. 2003), or As^{3+} -schwertmannite (Duquesne et al. 2003). *Thiomonas* strains, which oxidize both As^{3+} and Fe^{2+} , rapidly form Fe^{3+} - As^{5+} amorphous hydroxysulfates with As:Fe mole ratios up to 0.7 (Leblanc et al. 1996; Morin et al. 2003). This value approaches the upper limits observed for synthetic coprecipitates of As^{5+} -hydrous ferric oxides (Carlson et al. 2002). EXAFS analysis of these mineral phases confirms the similarity between the natural and *in vitro* mineral phases. In addition, spectroscopic evidence of the formation of stable arsenic complexes on the schwertmannite surface (Waychunas et al. 1995) explains the inhibiting role of As in the crystallization of schwertmannite, leading to the formation of amorphous As-rich ferric hydroxysulfates (Carlson et al. 2002).

Eventually, farther downstream in the AMD, the mixing of acid waters from former mining sites, with typical pH values of 2–3, with non-polluted neutral waters strongly modifies arsenic speciation (Gault et al. 2003; Casiot et al. 2005). Neutralization at the confluence leads to a solubilization of arsenate and arsenite complexes and to the oxidation of any residual Fe^{2+} . Most of the remaining As is sorbed at the surface of the precipitated hydrous Fe- and Al-oxides, which settle out during the first hundreds of meters downstream. Traces of arsenic will be transported farther in colloidal form (Kimball et al. 1995; Casiot et al. 2005).

ARSENIC SPECIATION IN SOILS

Arsenic can accumulate in soils (Voigt et al. 1996), contaminate both surface waters and groundwaters, and be taken up by plants and animals, thus affecting the whole food chain. The arsenic may be derived from the parent rocks or introduced anthropogenically by industrial activities, such as the mining and processing of arsenic or the manufacture

of arsenic chemicals, or by agriculture through various pesticides. The global average concentration of arsenic in uncontaminated soil is 5 to 6 mg kg^{-1} , with variations of more than an order of magnitude, depending on the kind of soil considered (Chen et al. 2002). Although the regulations governing As contamination in waters are well defined, the regulatory goals for remediation of contaminated soils vary greatly among countries and even locally, and are around 1 to 25 mg kg^{-1} for residential soils. Important issues centre on the bioavailability of the arsenic, which is related to its speciation, as well as the change of As mobility (and bioavailability) with time.

Iron oxides play a pivotal role in the biogeochemical behavior of As. They provide a ubiquitous As-trapping process in environments such as sediments (Belzile and Tessier 1990) and soils (Morin et al. 2002; Cancès et al. 2005). Hydrous ferric oxides (HFO), e.g. ferrihydrite, are the most reactive soil components as regards As sorption and can take up hundreds of mg kg^{-1} As, either as As^{3+} or As^{5+} (Goldberg 2002 and references therein). Acidic conditions favor the protonation of mineral surfaces and the adsorption of arsenate polyanions, H_2AsO_4^- or HAsO_4^{2-} , while As^{3+} remains soluble in such conditions, as arsenious acid, H_3AsO_3 . Under alkaline conditions, As^{3+} adsorbs preferentially to As^{5+} on iron oxide surfaces. Near neutral pH, i.e. closer to the *pKa* of H_3AsO_3 (9.2), arsenate and arsenite both adsorb with the same efficiency at the surfaces of HFO and crystalline iron oxides via the formation of strong inner-sphere surface complexes. Similar bidentate binuclear sorption mechanisms have been found for As^{3+} (Manning et al. 1998; Farquhar et al. 2002; Ona-Nguema et al. 2005) and As^{5+} (Waychunas et al. 1996; Sherman and Randall 2003), although there is evidence for an additional bidentate mononuclear complex for As^{3+} , especially in the case of hematite and ferrihydrite (Ona-Nguema et al. 2005).

Anoxic conditions, e.g. in the saturated zone of hydromorphic soils or in groundwater, may favor the reduction of Fe^{3+} and As^{5+} , often assisted by bacterial activity involving dissimilatory iron-reducing bacteria (DIRB) and dissimilatory arsenic-reducing bacteria (DARB) (see Lloyd and Oremland this issue). In natural environments, iron reduction is generally accompanied by the reduction of As, which can lead to the release of soluble As^{3+} . However, the respective roles of the reduction of iron and arsenic in the mobilization of arsenic are still not clear (van Geen et al. 2004), and in this context, some specific characteristics of As^{3+}




FIGURE 2 Changes in As speciation in a soil profile (A) developed on mineralized schists in the Echassières region, France. (B) As, Cu, and Zn concentrations indicate some enrichment relative to the bedrock in the upper part of the saprolite (B and C₁ horizons) and a progressive depletion towards surficial A₂–A₀₀ horizons. (C) XRD patterns show the progressive disappearance of primary pharmacosiderite (Ph) towards the upper part of the saprolite; the main Bragg reflections of muscovite (Mu) and biotite (Bt) are also indicated. (D) Fourier filtered contributions of the second neighbor contributions to the EXAFS spectra at As K-edge indicating changes in As speciation between the saprolite and the upper horizons; the spectra are similar to those of pharmacosiderite and As adsorbed on HFO, respectively. (E) SEM photograph displaying the typical euhedral cubic crystals of pharmacosiderite formed after hydrothermal weathering of primary iron–arsenic sulfide minerals. Modified from Morin et al. (2002)

speciation need to be taken into account, especially its low sorption efficiency on clay and Al-hydroxide surfaces as compared to As⁵⁺ (Goldberg 2002).

We discuss here two contrasting examples of arsenic-rich soils, one a former industrial site and the other a natural “geochemical anomaly.” Both emphasize the role of iron oxides in the immobilization of arsenic in soils. The first example is a former pesticide plant at Auzon (Haute-Loire, France). Here, industrial activities started at the beginning of the 20th century and lasted until 1949. The first activity at this site was the roasting of arsenic sulfides [arsenopyrite and realgar (As₂S₃)] from the neighboring mines in order to produce arsenolite (As₂O₃). Subsequently, other As-containing chemical products were manufactured at the site, including arsenical pesticides. The high arsenic concentrations encountered at this site (up to 1.2 wt% in topsoils and 35 mg L⁻¹ in groundwaters) have motivated research to understand the processes of contamination of the soils and underlying sediments (Cancès et al. 2005). The upper horizons contain arsenic minerals inherited from the contaminated site, such as arseniosiderite, formed by oxidation of sulfide ores, and schultenite (PbHAsO₄), manufactured for pesticides. However, detailed analysis of arsenic speciation throughout several soil profiles revealed that most of the arsenic was present as arsenate scavenged by Fe-oxyhydroxides. However, detailed analysis of arsenic speciation throughout several soil profiles revealed that most of the arsenic was present as arsenate scavenged by Fe-oxyhydroxides. This trapping process appeared to take place after the breakdown of source minerals such as schultenite.

Geochemical anomalies provide examples of regional, long-term environmental contamination. At Echassières (Allier, France) in the Massif Central (Fig. 2; Morin et al. 2002), soils have developed on schists enriched in As over tens of km². This has led to As concentrations of up to 5000 mg kg⁻¹ in the lowermost weathered horizon (saprolite) and up to 900 mg kg⁻¹ at the soil surface. The primary As-bearing minerals in the underlying rocks are arsenopyrite, löllingite (FeAs₂), and a pharmacosiderite solid solution close to the barium-pharmacosiderite end-member, [BaFe₄(AsO₄)₃(OH)₅•5H₂O]. This pharmacosiderite formed during a late hydrothermal event at the expense of sulfides and arsenides and represents 70% of the total As in the saprolite. However, it accounts for only 20–30% of total As in the upper horizons, which display a progressive decrease in As concentration upwards through the

soil profile. The rest of the arsenic in the soils occurs as arsenate adsorbed as inner-sphere complexes at the surfaces of poorly crystalline Fe-oxides. In this case, adsorption provides long-term trapping of the arsenic, given slightly acidic and oxidizing conditions, and delays potential release of As into the biosphere.

Finally, the release of arsenic through agricultural activities remains a great concern. Various approaches have been tried to reduce the leachability of arsenic from pesticides in order to protect surface water and groundwater from contamination and reduce the bioavailability of arsenic to human receptors. Adding Fe- and Mn-containing materials to cause adsorption and coprecipitation of the arsenic has been one remediation strategy. On the other hand, addition of phosphate-based components increases the leachability of arsenic, whereas mixed phosphate–iron additions reduce both the leachability and the bioaccessibility of arsenic relative to the unamended soils (Martin and Ruby 2003). Much remains to be done in order to develop a clear picture of the complex interplay among the impacts of As on the biosphere, on soil biogeochemistry, and on remediation technologies.

ACKNOWLEDGMENTS

We thank Guest Editor David Vaughan for the invitation to prepare this review article. This work has been made possible over the years through fruitful collaborations with F. Juillot and our late colleague P. Ildefonse at IMPMC (Paris, France), M. Benedetti (LGM, Paris, France), F. Elbaz-Poulichet and her group (Hydrosciences, Montpellier, France), V. Bonnefoy (LCM, Marseille, France), G. E. Brown Jr. (Department of Earth and Environmental Sciences, Stanford University, USA), and the members of the Stanford Environmental Molecular Science Institute (NSF-SEMSI). The authors are indebted to V. Briois at LURE (Orsay, France), J.-L. Hazemann at FAME/ESRF (Grenoble, France), and J. Bargar at SSRL (Stanford, USA) for their technical assistance during synchrotron experiments. The manuscript has been greatly improved by editors David Vaughan, Mike Hochella and Pierrette Tremblay. This work was supported by CNRS (PNSE, GEOMEX, and ECCO programs). This is IPGP contribution #2122. ■

REFERENCES

- Alpers CN, Jambor JL, Nordstrom DK (2000) Sulfate Minerals. Crystallography, Geochemistry and Environmental Significance. Reviews in Mineralogy and Geochemistry 40, Mineralogical Society of America and Geochemical Society
- Anthony JW, Bideaux RA, Bladh KW, Nichols MC (2000) Handbook of Mineralogy, Volume IV: Arsenates, Phosphates, Vanadates. Mineral Data Publishing, Tucson, USA
- Belzile N, Tessier A (1990) Interactions between arsenic and iron oxyhydroxides in lacustrine sediments. *Geochimica et Cosmochimica Acta* 54: 103-109
- Bigham JM, Schwertmann U, Traina SJ, Winland RL, Wolf M (1996) Schwertmannite and the chemical modeling of iron in acid sulfate waters. *Geochimica et Cosmochimica Acta* 60: 2111-2121
- Brown GE Jr, Foster AL, Ostergren JD (1999) Mineral surfaces and bioavailability of heavy metals: A molecular-scale perspective. Proceedings of the National Academy of Sciences USA 96: 3388-3395
- Bruneel O, Personné J-C, Casiot C, Leblanc M, Elbaz-Poulichet F, Mahler BJ, Le Flèche A, Grimont PAD (2003) Mediation of arsenic oxidation by *Thiomonas* sp. in acid mine drainage (Carnoulès, France). *Journal of Applied Microbiology* 95: 492-499
- Cancès B, Juillot F, Morin G, Laperche V, Alvarez L, Proux O, Hazemann J-L, Brown GE Jr, Calas G (2005) XAS evidence of As(V) association with iron oxyhydroxides in a contaminated soil at a former arsenical pesticide processing plant. *Environmental Science & Technology* 39: 9398-9405
- Carlson L, Bigham JM, Schwertmann U, Kyek A, Wagner F (2002) Scavenging of As from acid mine drainage by schwertmannite and ferrihydrite: A comparison with synthetic analogues. *Environmental Science & Technology* 36: 1712-1719
- Casiot C, Lebrun S, Morin G, Bruneel O, Personné JC, Elbaz-Poulichet F (2005) Sorption and redox processes controlling arsenic fate and transport in a stream impacted by acid mine drainage. *Science of the Total Environment* 347: 122-130
- Chen M, Ma LQ, Harris WG (2002) Arsenic concentrations in Florida surface soils: Influence of soil type and properties. *Soil Science Society of America Journal* 66: 632-640
- Cornell RM, Schwertmann U (2003) The iron oxides: structure properties, reactions, occurrence and uses. VCH Weinheim, Germany
- Courtin-Nomade A, Grosbois C, Bril H, Roussel C (2005) Spatial variability of arsenic in some iron-rich deposits generated by acid mine drainage. *Applied Geochemistry* 20: 383-396
- Duker AA, Carranza EJM, Hale M (2005) Arsenic geochemistry and health. *Environment International* 31: 631-641
- Duquesne K, Lebrun S, Casiot C, Bruneel O, Personné J-C, Leblanc M, Elbaz-Poulichet F, Morin G, Bonnefoy V (2003) Immobilization of arsenite and ferric iron by *Acidithiobacillus ferrooxidans* and its relevance to acid mine drainage. *Applied and Environmental Microbiology* 69: 6165-6173
- Farquhar ML, Charnock JM, Livens FR, Vaughan DJ (2002) Mechanisms of arsenic uptake from aqueous solution by interaction with goethite, lepidocrocite, mackinawite, and pyrite: An X-ray absorption spectroscopy study. *Environmental Science & Technology* 36: 1757-1762
- Foster AL, Brown Jr GE, Tingle TN, Parks GA (1998) Quantitative arsenic speciation in mine tailings using X-ray absorption spectroscopy. *American Mineralogist* 83: 553-568
- Fukushi K, Sasaki M, Sato T, Yanase N, Amano H, Ikeda H (2003) A natural attenuation of arsenic in drainage from an abandoned arsenic mine dump. *Applied Geochemistry* 18: 1267-1278
- Gault AG, Polya DA, Lythgoe PR, Farquhar ML, Charnock JM, Wogelius RA (2003) Arsenic speciation in surface waters and sediments in a contaminated waterway: an IC-ICP-MS and XAS based study. *Applied Geochemistry* 18: 1387-1397
- Goldberg S (2002) Competitive adsorption of arsenate and arsenite on oxide and clay minerals. *Soil Science Society of America Journal* 66:413-421
- Inskeep WP, Macur RE, Harrison G, Bostick BC, Fendorf S (2004) Biomineralization of As(V)-hydrous ferric oxyhydroxide in microbial mats of an acid-sulfate-chloride geothermal spring, Yellowstone National Park. *Geochimica et Cosmochimica Acta* 68: 3141-3155
- Juillot F, Ildefonse P, Morin G, Calas G, De Kersabiec AM, Benedetti M (1999) Remobilization of arsenic from buried wastes in an industrial site: mineralogical and geochemical control. *Applied Geochemistry* 14: 1031-1048
- Kimball BA, Callender E, Axtmann EV (1995) Effects of colloids on metal transport in a river receiving acid mine drainage, upper Arkansas River, Colorado, U.S.A. *Applied Geochemistry* 10: 285-306
- Leblanc M, Achard B, Ben Othman D, Luck JM, Bertrand-Sarfati J, Personné J-C (1996) Accumulation of arsenic from acidic mine waters by ferruginous bacterial accretions (stromatolites). *Applied Geochemistry* 11: 541-549
- Manning BA, Fendorf SE, Goldberg S (1998) Surface structures and stability of arsenic(III) on goethite: spectroscopic evidence for inner-sphere complexes. *Environmental Science & Technology* 32: 2383-2388
- Martin TA, Ruby MV (2003) In situ remediation of arsenic in contaminated soils. *Remediation Journal* 14: 21-32
- Moon DH, Dermatas D, Menounou N (2004) Arsenic immobilization by calcium-arsenic precipitates in lime treated soils. *Science of the Total Environment* 330: 171-185
- Morin G, Lecocq D, Juillot F, Calas G, Ildefonse P and eight authors (2002) EXAFS evidence of sorbed arsenic(V) and pharmacosiderite in a soil overlying the Echassières geochemical anomaly, Allier, France. *Bulletin de la Société Géologique de France* 173: 281-291
- Morin G, Juillot F, Casiot C, Bruneel O, Personné J-C, Elbaz-Poulichet F, Leblanc M, Ildefonse P, Calas G (2003) Bacterial formation of tooleite and mixed arsenic(III) or arsenic(V)-iron(III) gels in the Carnoulès acid mine drainage. A XANES, XRD, and SEM study. *Environmental Science & Technology* 37: 1705-1712
- Ona-Nguema G, Morin G, Juillot F, Brown GE Jr, Calas G (2005) EXAFS analysis of Arsenic(III) sorption onto 2-line ferrihydrite, hematite, goethite, and lepidocrocite under anoxic conditions. Influence of the surface structure. *Environmental Science and Technology* 39, 9147-9155
- Paktunc D, Foster A, Heald S, Laflamme G (2004) Speciation and characterization of arsenic in gold ores and cyanidation tailings using X-ray absorption spectroscopy. *Geochimica et Cosmochimica Acta* 68: 969-983
- Savage KS, Tingle TN, O'Day PA, Waychunas GA, Bird DK (2000) Arsenic speciation in pyrite and secondary weathering phases, Mother Lode Gold District, Tuolumne County, California. *Applied Geochemistry* 15: 1219-1244
- Sherman DM, Randall SR (2003) Surface complexation of arsenic(V) to iron(III) oxides and oxide hydroxides: Structural mechanism from ab initio molecular geometries and EXAFS spectroscopy. *Geochimica et Cosmochimica Acta* 67: 4223-4230
- Swash PM, Monhemius A (1994) Hydrothermal precipitation from aqueous solutions containing iron(III) arsenate and sulphate. In: *Hydrometallurgy '94*. Chapman and Hall, New York, pp 177-190
- van Geen A, Rose J, Thorar S, Garnier JM, Zheng Y, Bottero JY (2004) Decoupling of As and Fe release to Bangladesh groundwater under reducing conditions. Part II: Evidence from sediment incubations. *Geochimica et Cosmochimica Acta* 68: 3475-3486
- Voigt DE, Brantley SL, Hennet RJC (1996) Chemical fixation of arsenic in contaminated soils. *Applied Geochemistry* 11: 633-643
- Waychunas GA, Xu N, Fuller CC, Davis JA, Bigham JM (1995) XAS study of AsO₄³⁻ and SeO₄²⁻ substituted schwertmannites. *Physica B* 208&209: 481-483
- Waychunas GA, Fuller CC, Rea BA, Davis JA (1996) Wide angle X-ray scattering (WAXS) study of "two-line" ferrihydrite structure: Effect of arsenate sorption and counterion variation and comparison with EXAFS results. *Geochimica et Cosmochimica Acta* 60: 1765-1781
- Williams M (2001) Arsenic in mine waters: an international study. *Environmental Geology* 40: 267-278

3rd International Workshop on Highly Siderophile Element Geochemistry

The 3rd international workshop on Highly Siderophile Element Geochemistry: High-Temperature and Low-Temperature Processes, Environment and Health Issues will be held at the Department of Earth Sciences, Durham University, UK, 5-7 July 2006. For details, see: www.dur.ac.uk/earth.sciences/conferences/ or e-mail highly.siderophile@durham.ac.uk

The Maine Pegmatite Workshop

Allison Brand

University of British Columbia, Canada

Thanks in part to a generous travel grant from the Mineralogical Association of Canada, I was able to attend the Maine Pegmatite Workshop (June 4–13, 2005) in Poland, Maine (USA). The workshop was led by Dr. William “Skip” Simmons and Dr. Karen Webber, from the University of New Orleans, and Raymond Sprague, Maine pegmatite miner and administrator. I couldn’t have imagined a more inspiring, intimate atmosphere at an international meeting. Researchers, miners, mineral collectors, students, rockhounds, artists, and general pegmatite enthusiasts came together from all parts of the globe, with open minds and valuable perspectives, to learn about and discuss the multidisciplinary world of pegmatite.

The 34 participants assembled each morning for lectures and discussions and headed out each afternoon to a different pegmatite quarry or mine. Western Maine is a well-known pegmatite district with a dense history of mining and exploration; some of the more highly evolved pegmatites were exploited for mica and feldspar during the late 1800s and early 1900s. Many have yielded excellent examples of gem tourmaline, purple apatite, and a lot more. The field trips included a full-day excursion to the Palermo mine in New Hampshire, by invitation of its owner, Bob Whitmore. Palermo is a world-famous site for phosphates and is a type locality for ten mineral species. Other days included trips to Mount Mica, famous for its amazing gem tourmalines and quartz crystals, and to BB7, the Hole-in-the Ground, Pulsifer, Bennet Mt., Marie, and Emmons quarries. Seeing the glow of satisfaction on the faces of participants when they had just discovered a perfect quartz crystal, a brilliant tourmaline, or a euhedral beryl were some of my favourite moments.

What ignited *my* interest was the stimulating morning discussion. Here was a forum where a diverse group with varied perspectives could debate the controversial issues, and each participant’s ideas were as valued as the next’s. It was exciting to watch researchers present their theories and experiments, which could be supported or questioned by what miners were seeing in the field. For example, large crystal size is usually associated with slow crystal growth. However, Dr. Webber’s research suggests that pegmatite bodies may crystallize in a matter of days or months given the right diffusion and nucleation conditions, much like large ice crystals can grow overnight. Some of the discussions became quite lively, depending on the level of controversy associated with the topic. How should we classify pegmatites, and what is the origin of the quartz core common in zoned pegmatites? The “A-word” (anatexis) was also discussed as a possible origin for pegmatites, a topic hotly debated internationally.



Group picture of the Maine Pegmatite Workshop 2005

With the aid of their book, *Pegmatology*, Dr. Simmons and Dr. Webber guided the group through the basics of pegmatite science, plate tectonics, the origin of pockets, and pocket indicators. Faculty included Al Forster and Jim Nizamoff, also from the University of New Orleans, Ray Sprague, and the original “Pocket Fairy” himself, Frank C. Perham, with whom I had the pleasure of carpooling every day to the field sites. Frank has been a pegmatite miner and expert for over 50 years and seems to have been present at every major pocket discovery in the district! Invited speakers took over for the later lectures and evening programs, and included a worldwide tour of pegmatite

localities (Mike Wise), and more specifically Pakistan (Brendan Laurs), Brazil (Odulio J.M. de Moura), the Northwest Territories of Canada (Lee Groat), the Himalaya Pegmatite in San Diego County, California (Jesse Fisher), the Karelia beryl deposit in Finland (Peter Lyckberg), and the alkaline pegmatites of Zhomba-Malosa in Malawi (Alessandro Guastani). Evening events also included a Maine lobster feast, a display of the prize specimens and wares of participants and local mineral collectors, an exhibition of the fine mineral art of award-winning artist Fred Wilda, and a never-to-be-forgotten “Ugly Shirt Contest.”

Pegmatites were first introduced to me by my undergraduate thesis supervisor, Prof. Lee Groat of UBC, with whom I have worked for three years. For my thesis, I studied pegmatite-related emerald mineralization in northwestern Ontario, but I had also been able to visit the Little Nahanni Pegmatite Group (NWT, Canada) with Elspeth Barnes (completing a PhD on the area) and Prof. Groat the previous summer. The latter area is very large, containing hundreds of pegmatite dikes, and offers almost complete 3-D exposure of the pegmatite field, with amazing textures and mineralization. In the open-minded environment of the Maine Pegmatite Workshop, I felt completely comfortable contributing my own limited experiences to the conversation, a confidence I have never felt in a previous scientific forum. As pegmatite science evolves and attempts to answer some of the more controversial and difficult questions, this annual workshop can only grow more exciting.

The fifth annual Maine Pegmatite Workshop will be held May 27–June 4, 2006
homepage.mac.com/rasprague/PegShop



Allison in the field in the Northwest Territories